

Toda-Darboux maps and vertex operators

M. Adler* P. van Moerbeke†

February 9, 2008

Consider the Toda lattice

$$\frac{\partial L}{\partial t_n} = [(L^n)_+, L] = [-(L^n)_-, L], \quad n = 1, 2, \dots \quad (0.1)$$

on bi-infinite tridiagonal matrices

$$L = \Lambda^{-1}a + b\Lambda^0 + \Lambda, \quad (0.2)$$

with Λ being the customary shift $\Lambda := (\delta_{i,j-1})_{i,j \geq 0}$ and a and b being diagonal matrices. As is well known, in analogy with Sato's KP-theory, the entries a and b have the following τ -function representation

$$b_n = \frac{\partial}{\partial t_1} \log \frac{\tau_{n+1}}{\tau_n} \quad \text{and} \quad a_{n-1} = \frac{\tau_{n-1}\tau_{n+1}}{\tau_n^2}, \quad (0.3)$$

in terms of a τ -vector $\tau = (\tau_n)_{n \in \mathbb{Z}}$. In section 2, it is shown that, if τ is a Toda lattice τ -vector, then

$$(1 + c \mathbb{X}(t, z))\tau$$

*Department of Mathematics, Brandeis University, Waltham, Mass 02254, USA. The support of a National Science Foundation grant # DMS-9503246 is gratefully acknowledged.

†Université de Louvain, 1348 Louvain-la-Neuve, Belgium and Department of Mathematics, Brandeis University, Waltham, Mass 02254, USA. The support of a National Science Foundation grant # DMS-9503246, a Nato, a FNRS and a Francqui Foundation grant is gratefully acknowledged.

is a Toda τ -vector as well, where the *vertex operator* $X(t, z)$, first introduced by us in [3], is given by¹

$$\mathbb{X}(t, z) = \Lambda^{-1} \chi(z^2) e^{\sum t_i z^i} e^{-2 \sum \frac{z^{-i}}{i} \frac{\partial}{\partial t_i}}. \quad (0.4)$$

For a fixed $\lambda \in \mathbb{C}^*$, we consider a Borel factorization

$$L(t) - \lambda = L_-(t) L_+(t), \text{ with } L_-(t) = \Lambda^{-1} \alpha(t) + \Lambda^0, \quad L_+(t) = \beta(t) \Lambda^0 + \Lambda$$

and the corresponding *Darboux transform*

$$L(t) - \lambda = L_- L_+ \mapsto \tilde{L}(t) - \lambda = L_+ L_-, \quad (0.5)$$

which in coordinates reads:

$$\begin{aligned} b_n - \lambda &= \alpha_{n-1} + \beta_n \mapsto \tilde{b}_n - \lambda = \alpha_n + \beta_n = (b_n - \lambda) + (\alpha_n - \alpha_{n-1}) \\ a_{n-1} &= \alpha_{n-1} \beta_{n-1} \mapsto \tilde{a}_{n-1} = \alpha_{n-1} \beta_n = a_{n-1} \frac{\beta_n}{\beta_{n-1}}. \end{aligned} \quad (0.6)$$

This map will be called *Toda-Darboux*, when both matrices $L - \lambda = L_- L_+$, and $\tilde{L} = L_+ L_-$ flow according to all Toda vector fields.

The following *vertex operators* will play an important role in this paper:

$$\mathbb{X}_1(t, \lambda) := \chi(\lambda) X(t, \lambda) \quad \text{and} \quad \mathbb{X}_2(t, \lambda) := \chi(\lambda^{-1}) X(-t, \lambda) \Lambda, \quad (0.7)$$

where

$$X(t, \lambda) := e^{\sum_1^\infty t_i \lambda^i} e^{-\sum_1^\infty \frac{\lambda^{-i}}{i} \frac{\partial}{\partial t_i}}. \quad (0.8)$$

We also define a *discrete Wronskian* $\{ , \}$ on column vectors f and g

$$\{f, g\} := (f_{n+1} g_n - f_n g_{n+1})_{n \in \mathbb{Z}}. \quad (0.9)$$

Theorem 0.1. *Each element of the 2-dimensional null-space*

$$\ker(L(t) - \lambda) = \left\{ \Phi(t, \lambda) = \frac{\tilde{\tau}(t)}{\tau(t)} = \frac{\left(a \mathbb{X}_1(t, \lambda) + b e^{\sum t_i \lambda^i} \mathbb{X}_2(t, \lambda) \right) \tau(t)}{\tau(t)} \right\}, \quad (0.10)$$

¹ $\chi(z) := \text{diag}(\dots, z^{-1}, 1, z, \dots)$

where $\Phi(t, \lambda)$ satisfies

$$\frac{\partial \Phi}{\partial t_n} = (L^n)_+ \Phi,$$

specifies a factorization for all $t \in \mathbb{C}^\infty$,

$$L(t) - \lambda = L_- L_+ = (\Lambda^{-1} \alpha + \Lambda^0)(\beta \Lambda^0 + \Lambda),$$

with

$$\alpha = \Lambda \frac{\partial}{\partial t_1} \log \Phi(t, \lambda) - \lambda \text{ and } \beta = -\frac{\Lambda \Phi(t, \lambda)}{\Phi(t, \lambda)}, \quad (0.11)$$

and so

$$\ker L_+ = \mathbb{C} \Phi(t, \lambda).$$

The Darboux transform (0.5) maps $L(t) - \lambda$ into a new tridiagonal matrix $\tilde{L}(t) - \lambda$, with entries $\tilde{b}_n$ and $\tilde{a}_n$ given by (0.3) in term of $\tilde{\tau}$. The Toda-Darboux transform on L induces a map on τ :

$$\tau \longmapsto \tilde{\tau} = \tau \Phi = \left(a \mathbb{X}_1(t, \lambda) + b e^{\sum t_i \lambda^i} \mathbb{X}_2(t, \lambda) \right) \tau(t). \quad (0.12)$$

The kernel $\ker(L(t) - \lambda)$, as in (0.10), contains two distinguished (wave) functions, whose asymptotics is given later in (2.5),

$$\Phi^{(1)}(t, \lambda) := \frac{\mathbb{X}_1(t, \lambda) \tau(t)}{\tau(t)} \text{ and } \Phi^{(2)}(t, \lambda) := \frac{e^{\sum t_i \lambda^i} \mathbb{X}_2(t, \lambda) \tau(t)}{\tau(t)} \quad (0.13)$$

which Darboux transform, as follows:

Theorem 0.2. *The wave functions $\Phi^{(1)}$ and $\Phi^{(2)}$ for the L -operator are Darboux transformed into wave functions $\tilde{\Phi}^{(1)}$ and $\tilde{\Phi}^{(2)}$ for the Darboux transformed operator $\tilde{L}$; they are given by Wronskian formulas, also expressible in terms of the new τ -function $\tilde{\tau}$, to wit*

$$\begin{aligned} \tilde{\Phi}^{(1)}(t, z) &= \frac{\mathbb{X}_1(t, z) \tilde{\tau}}{\tilde{\tau}} = \frac{1}{z} \frac{\{\Phi^{(1)}(t, z), \Phi(t, \lambda)\}}{\Phi(t, \lambda)} = \frac{1}{z} L_+ \Phi^{(1)}(t, z) \\ \tilde{\Phi}^{(2)}(t, z) &= \frac{e^{\xi(t, z)} \mathbb{X}_2(t, z) \tilde{\tau}}{\tilde{\tau}} = \frac{z}{\lambda - z} \frac{\{\Phi^{(2)}(t, z), \Phi(t, \lambda)\}}{\Phi(t, \lambda)} = \frac{z}{\lambda - z} L_+ \Phi^{(2)}(t, z), \end{aligned}$$

thus satisfying

$$\tilde{L}\tilde{\Phi}^{(i)} = z\tilde{\Phi}^{(i)} \text{ and } \frac{\partial \tilde{\Phi}^{(i)}}{\partial t_n} = (\tilde{L}^n)_+ \tilde{\Phi}^{(i)}, \quad i = 1, 2.$$

Theorem 0.3. *The following holds:*

$$\text{Toda for } L \text{ and } \tilde{L} \iff \text{KM-lattice for } L_- \text{ and } L_{+,}.$$

In coordinates, the latter takes the form

$$\dot{\alpha}_n = (\beta_{n+1} - \beta_n)\alpha_n, \quad \dot{\beta}_n = (\alpha_n - \alpha_{n-1})\beta_n; \quad (0.14)$$

moreover α_n and β_n satisfy the Ricatti equations, with coefficients given by the entries of L ,

$$\begin{aligned} \dot{\alpha}_n &= -\alpha_n^2 + (b_{n+1} - \lambda)\alpha_n - a_n \\ \dot{\beta}_n &= \beta_n^2 - (b_n - \lambda)\beta_n + a_n. \end{aligned}$$

in well known analogy with the Sturm-Liouville situation.

Consider *band matrices* of the form

$$L = \sum_{-p \leq i \leq p} a_i \Lambda^i, \quad a_p = I, \quad (0.15)$$

with a_i being diagonal matrices; define Toda lattice vector fields, as follows:

$$\begin{aligned} \frac{\partial L}{\partial x_i} &= [(\overline{L^{i/p}})_+, L], \quad \frac{\partial L}{\partial y_i} = [(L^{i/p})_-, L], \text{ for } i = 1, 2, \dots, p \not\equiv i \\ \frac{\partial L}{\partial t_{ip}} &= [(L^i)_+, L], \quad i = 1, 2, \dots \end{aligned} \quad (0.16)$$

Note $\overline{L^{i/p}}$ and $\underline{L^{i/p}}$ involve *right* p^{th} roots and *left* p^{th} roots:

$$\begin{aligned} \overline{L^{i/p}} &= (\overline{L^{1/p}})^i = \left(\Lambda + \sum_{k \leq 0} b_k \Lambda^k \right)^i \\ \underline{L^{i/p}} &= (\underline{L^{1/p}})^i = \left(c_{-1} \Lambda^{-1} + \sum_{k \geq 0} c_k \Lambda^k \right)^i; \end{aligned} \quad (0.17)$$

with this notation, the vector fields (0.16) preserve the band structure of L .

Then L can be expressed in terms of a string of τ -functions,

$$\tau_n := \tau_n(\hat{x}, \hat{y}, \hat{t}), \quad (0.18)$$

with $\hat{x}$, $\hat{y}$, and $\hat{t}$ having certain components omitted:

$$\hat{x} = (x_1, \dots, \hat{x}_p, \dots, \hat{x}_{2p}, \dots), \quad \hat{y} = (y_1, \dots, \hat{y}_p, \dots, \hat{y}_{2p}, \dots), \quad \hat{t} = (t_p, t_{2p}, t_{3p}, \dots). \quad (0.19)$$

Define the following vertex operators:

$$\begin{aligned} \mathbb{X}_1(\lambda) &= \chi(\lambda) e^{\sum_1^\infty t_{pi} \lambda^{pi}} e^{-\sum_1^\infty \frac{\lambda^{-pi}}{pi} \frac{\partial}{\partial t_{pi}}} e^{\sum_{p \nmid i} x_i \lambda^i} e^{-\sum_{p \nmid i} \frac{\lambda^{-i}}{i} \frac{\partial}{\partial x_i}} \\ \mathbb{X}_2(\lambda) &= \chi(\lambda^{-1}) e^{-\sum_1^\infty t_{pi} \lambda^{pi}} e^{\sum_1^\infty \frac{\lambda^{-pi}}{pi} \frac{\partial}{\partial t_{pi}}} e^{\sum_{p \nmid i} y_i \lambda^i} e^{-\sum_{p \nmid i} \frac{\lambda^{-i}}{i} \frac{\partial}{\partial y_i}}. \end{aligned}$$

Theorem 0.4. *Each element of the $2p$ -dimensional null-space²*

$$\ker(L - \lambda^p) = \left\{ \Phi(\lambda) = \frac{\tilde{\tau}}{\tau} = \frac{\sum_{k=0}^{p-1} \left(a_k \mathbb{X}_1(\omega^k \lambda) + b_k e^{\sum_1^\infty t_{ip} \lambda^{ip}} \mathbb{X}_2(\omega^k \lambda) \right) \tau}{\tau} \right\},$$

where $\Phi(\lambda)$ satisfies

$$\begin{aligned} L\Phi &= \lambda^p \Phi \\ \frac{\partial \Phi}{\partial x_i} &= (L^{i/p})_+ \Phi, \quad \frac{\partial \Phi}{\partial y_i} = (L^{i/p})_- \Phi, \quad \text{for } i = 1, 2, \dots \text{ with } p \nmid i \\ \frac{\partial \Phi}{\partial t_{ip}} &= (L^i)_+ \Phi, \quad \text{for } i = 1, 2, \dots, \end{aligned} \quad (0.20)$$

determines a Toda-Darboux transform

$$L - \lambda^p \longmapsto (\beta \Lambda^0 + \Lambda)(L - \lambda^p)(\beta \Lambda^0 + \Lambda)^{-1}$$

with

$$\beta = -\frac{\Lambda \Phi(\lambda)}{\Phi(\lambda)};$$

² ω is a primitive p th root of unity.

it acts on τ as

$$\tau \longmapsto \tilde{\tau} = \tau \Phi = \sum_{k=0}^{p-1} \left(a_k BX_1(\omega^k \lambda) + b_k e^{\sum_{i=1}^{\infty} t_{ip} \lambda^{ip}} \mathbb{X}_2(\omega^k \lambda) \right) \tau.$$

For a broad account of Darboux transforms in a variety of situations, see the book of Matveev and Salle [9], which also contains a very extensive bibliography. Darboux transforms for differential operators, and their connections with τ -functions have been investigated in [8, 2, 12] for 2nd order differential operators and by Bakalov, Horozov and Yakimov [6] for general differential operators. In fact, using our vertex operator methods (see [2]), the latter results can be made quite a bit more precise. The connection with the KM-lattice was first made in [1].

Table of contents:

1. The 2-Toda lattice
2. Reduction from 2-Toda to tridiagonal 1-Toda, and vertex operators
3. Toda-Darboux transformations
4. Wronskians and Vertex operators for 1-Toda
5. Borel factorization, KM-lattice and Ricatti equations
6. Toda flows and Darboux transforms for band matrices

1 The 2-Toda lattice

Consider the splitting of the algebra $\mathcal{D}$ of pairs (P_1, P_2) of infinite $(\mathbb{Z} \times \mathbb{Z})$ matrices such that $(P_1)_{ij} = 0$ for $j - i \gg 0$ and $(P_2)_{ij} = 0$ for $i - j \gg 0$, used in [5]; to wit:

$$\begin{aligned} \mathcal{D} &= \mathcal{D}_u + \mathcal{D}_\ell, \\ \mathcal{D}_u &= \{(P, P) \mid P_{ij} = 0 \text{ if } |i - j| \gg 0\} = \{(P_1, P_2) \in \mathcal{D} \mid P_1 = P_2\}, \\ \mathcal{D}_\ell &= \{(P_1, P_2) \mid (P_1)_{ij} = 0 \text{ if } j \geq i, (P_2)_{ij} = 0 \text{ if } i > j\}, \end{aligned}$$

with $(P_1, P_2) = (P_1, P_2)_u + (P_1, P_2)_\ell$ given by

$$\begin{aligned} (P_1, P_2)_u &= (P_{1+} + P_{2-}, P_{1+} + P_{2-}), \\ (P_1, P_2)_\ell &= (P_{1-} - P_{2-}, P_{2+} - P_{1+}); \end{aligned} \quad (1.1)$$

P_+ and P_- denote the upper (including diagonal) and strictly lower triangular parts of the matrix P , respectively.

The two-dimensional Toda lattice equations

$$\frac{\partial L}{\partial x_n} = [(L_1^n, 0)_u, L] \quad \text{and} \quad \frac{\partial L}{\partial y_n} = [(0, L_2^n)_u, L] \quad n = 1, 2, \dots \quad (1.2)$$

are deformations of a pair of infinite matrices

$$L = (L_1, L_2) = \left(\sum_{-\infty < i \leq 1} a_i^{(1)} \Lambda^i, \sum_{-1 \leq i < \infty} a_i^{(2)} \Lambda^i \right) \in \mathcal{D}, \quad (1.3)$$

where $a_i^{(1)}$ and $a_i^{(2)}$ are diagonal matrices depending on $x = (x_1, x_2, \dots)$ and $y = (y_1, y_2, \dots)$, such that

$$a_1^{(1)} = I \quad \text{and} \quad (a_{-1}^{(2)})_{nn} \neq 0 \quad \text{for all } n.$$

In their 2-Toda theory, Ueno-Takasaki [11] also introduce a pair of wave vectors $\Psi = (\Psi_1, \Psi_2)$, satisfying $(L_1, L_2)\Psi = (z, z^{-1})\Psi$ and³

$$\begin{cases} \frac{\partial}{\partial x_n} \Psi = (L_1^n, 0)_u \Psi = ((L_1^n)_+, (L_1^n)_+) \Psi \\ \frac{\partial}{\partial y_n} \Psi = (0, L_2^n)_u \Psi = ((L_2^n)_-, (L_2^n)_-) \Psi \end{cases} \quad (1.4)$$

In [11], it is shown that the wave vectors Ψ can be expressed in terms of one sequence of τ -functions $\tau(n, t, s) = \tau_n(t_1, t_2, \dots; s_1, s_2, \dots)$, $n \in \mathbb{Z}$, to wit:

$$\Psi_1(x, y; z) = \left(\frac{\tau_n(x - [z^{-1}], y)}{\tau_n(x, y)} e^{\sum_{i=1}^{\infty} x_i z^i} z^n \right)_{n \in \mathbb{Z}}, \quad (1.5)$$

$$\Psi_2(x, y; z) = \left(\frac{\tau_{n+1}(x, y - [z])}{\tau_n(x, y)} e^{\sum_{i=1}^{\infty} y_i z^{-i}} z^n \right)_{n \in \mathbb{Z}}, \quad (1.6)$$

³Here the action is viewed componentwise, e.g., $(A, B)\Psi = (A\Psi_1, B\Psi_2)$ or $(z, z^{-1})\Psi = (z\Psi_1, z^{-1}\Psi_2)$.

with τ satisfying the following bilinear identities:

$$\begin{aligned} & \oint_{z=\infty} \tau_n(x - [z^{-1}], y) \tau_{m+1}(x' + [z^{-1}], y') e^{\sum_{i=1}^{\infty} (x_i - x'_i) z^i} z^{n-m-1} dz \\ &= \oint_{z=0} \tau_{n+1}(x, y - [z]) \tau_m(x', y' + [z]) e^{\sum_{i=1}^{\infty} (y_i - y'_i) z^{-i}} z^{n-m-1} dz. \end{aligned} \quad (1.7)$$

for all $m, n \in \mathbb{Z}$. Conversely, any τ -vector satisfying the bilinear identity (1.7) leads to a solution of the 2-Toda lattice; see [10].

Upon introducing appropriate shifts of x and y , and evaluating the contour integration over a contour about ∞ and the singularities, created by the shifts, one finds the following Fay identities, due to [4]; they will be useful later:

$$\begin{aligned} & \tau_n(x - [z^{-1}], y + [v] - [u]) \tau_n(x, y) - \tau_n(x, y + [v] - [u]) \tau_n(x - [z^{-1}], y) \\ &= \frac{v - u}{z} \tau_{n+1}(x, y - [u]) \tau_{n-1}(x - [z^{-1}], y + [v]) \end{aligned} \quad (1.8)$$

and

$$\begin{aligned} & \tau_n(x, y + [v_1]) \tau_{n+1}(x + [z_1^{-1}] - [z_2^{-1}], y - [v_2]) \frac{z_1^{-1}}{z_1^{-1} - z_2^{-1}} \\ &+ \tau_n(x + [z_1^{-1}] - [z_2^{-1}], y + [v_1]) \tau_{n+1}(x, y - [v_2]) \frac{z_2^{-1}}{z_2^{-1} - z_1^{-1}} \\ &= \tau_{n+1}(x + [z_1^{-1}], y) \tau_n(x - [z_2^{-1}], y + [v_1] - [v_2]) \frac{v_1}{v_1 - v_2} \\ &+ \tau_{n+1}(x + [z_1^{-1}], y + [v_1] - [v_2]) \tau_n(x - [z_2^{-1}], y) \frac{v_2}{v_2 - v_1}. \end{aligned} \quad (1.9)$$

Consider the following 2-Toda vertex operators⁴

$$\mathbb{Y}_1(\lambda) = \chi(\lambda) X(x; \lambda) \quad \text{and} \quad \mathbb{Y}_2(\mu) = \chi(\mu) X(y, \mu^{-1}) \Lambda,$$

where $X(x, \lambda)$ is given by (0.8). In [4], we have shown that for fixed $\lambda, \mu \in \mathbb{C}$, the vertex operator

$$a \mathbb{Y}_1(\lambda) + b \mathbb{Y}_2(\mu)$$

maps 2-Toda τ -vectors into themselves. Spelled out,

$${}^4 \chi(\lambda) = (\lambda^n)_{n \in \mathbb{Z}}$$

$$\begin{aligned}
 & \left(\left(a\mathbb{Y}_1(\lambda) + b\mathbb{Y}_2(\mu) \right) \tau \right)_n \\
 &= ae^{\sum_1^\infty x_i \lambda^i} \lambda^n \tau_n(x - [\lambda^{-1}], y) + be^{\sum_1^\infty y_i \mu^{-i}} \mu^n \tau_{n+1}(x, y - [\mu]) \quad (1.10)
 \end{aligned}$$

is a new τ -vector for the 2-Toda lattice.

2 Reduction from 2- to tridiagonal 1-Toda, and vertex operators

In the notation of section 1, consider the locus of (L_1, L_2) 's in $\mathcal{D}_u$, i.e., such that $L_1 = L_2$. Since the equations (1.2) imply

$$\frac{\partial L_i}{\partial x_n} = [(L_1^n)_+, L_i] \quad \text{and} \quad \frac{\partial L_i}{\partial y_n} = [(L_2^n)_-, L_i] \quad n = 1, 2, \dots \quad (2.1)$$

we have that along $\mathcal{D}_u$

$$\frac{\partial(L_1 - L_2)}{\partial x_n} = 0 \quad \frac{\partial(L_1 - L_2)}{\partial y_n} = 0, \quad (2.2)$$

implying that the vector fields $\frac{\partial}{\partial t_n}$ and $\frac{\partial}{\partial s_n}$ are tangent to $\mathcal{D}_u$. Also when $L_1 = L_2$, the matrix $L := L_1 = L_2$ is tridiagonal. Moreover

$$\left(\frac{\partial}{\partial x_n} + \frac{\partial}{\partial y_n} \right) L_1 = [(L_1^n)_+ + (L_2^n)_-, L_1] = [(L_1^n)_+ + (L_1^n)_-, L_1] = 0; \quad (2.3)$$

setting

$$x_n = \frac{t_n + s_n}{2}, \quad y_n = \frac{-t_n + s_n}{2} \quad \text{and} \quad t_n = x_n - y_n, \quad s_n = x_n + y_n,$$

with

$$\frac{\partial}{\partial t_n} = \frac{1}{2} \left(\frac{\partial}{\partial x_n} - \frac{\partial}{\partial y_n} \right), \quad \frac{\partial}{\partial s_n} = \frac{1}{2} \left(\frac{\partial}{\partial x_n} + \frac{\partial}{\partial y_n} \right), \quad (2.4)$$

equation (2.1) implies that $L = L_1 = L_2$ is independent of s .

The 2-Toda lattice wave functions Ψ_1 and Ψ_2 , properly reduced, yield two distinguished eigenfunctions for the 1-Toda lattice; they have the following expressions in terms of the 1-Toda τ -functions:

$$\begin{aligned}
 \Phi^{(1)}(t, z) &:= \Psi_1(x, y; z) e^{-\sum_1^\infty y_i z^i} \\
 &= e^{\sum_1^\infty t_i z^i} \left(z^n \frac{\tau_n(t - [z^{-1}])}{\tau_n(t)} \right)_{n \in \mathbb{Z}} \\
 &= \frac{\mathbb{X}_1(z) \tau}{\tau} \\
 &= e^{\sum_1^\infty t_i z^i} z^n (1 + O(z^{-1})) \tag{2.5}
 \end{aligned}$$

$$\begin{aligned}
 \Phi^{(2)}(t, z) &:= \Psi_2(x, y; z^{-1}) e^{-\sum_1^\infty y_i z^i} \\
 &= \left(z^{-n} \frac{\tau_{n+1}(t + [z^{-1}])}{\tau_n(t)} \right)_{n \in \mathbb{Z}} \\
 &= \frac{e^{\sum_1^\infty t_i z^i} \mathbb{X}_2(z) \tau}{\tau} \\
 &= z^{-n} \left(\frac{\tau_{n+1}(t)}{\tau_n(t)} + O(z^{-1}) \right), \tag{2.6}
 \end{aligned}$$

in terms of the vertex operators, already defined in (0.7), namely

$$\mathbb{X}_1(z) = \chi(z) X(t, z) \quad \text{and} \quad \mathbb{X}_2(z) = \chi(z^{-1}) X(-t, z) \Lambda; \tag{2.7}$$

also, for later use, recall from (0.4),

$$\mathbb{X}(t, z) := \Lambda^{-1} \chi(z^2) e^{\sum t_i z^i} e^{-2 \sum \frac{z^{-i}}{i} \frac{\partial}{\partial t_i}}; \tag{2.8}$$

Using (1.4), (2.2) and the fact that $L_1 = L_2$, one checks

$$L\Phi^{(1)} = L_1\Phi^{(1)} = z\Phi^{(1)} \quad L\Phi^{(2)} = L_2\Phi^{(2)} = z\Phi^{(2)}$$

and

$$\frac{\partial \Phi^{(1)}}{\partial t_n} = \frac{1}{2} \left(\frac{\partial}{\partial x_n} - \frac{\partial}{\partial y_n} \right) \Psi_1(x, y; z) e^{-\sum_1^\infty y_i z^i} = (L^n)_+ \Phi^{(1)} \tag{2.9}$$

$$\frac{\partial \Phi^{(2)}}{\partial t_n} = \frac{1}{2} \left(\frac{\partial}{\partial x_n} - \frac{\partial}{\partial y_n} \right) \Psi_2(x, y; z^{-1}) e^{-\sum y_i z^i} = (L^n)_+ \Phi^{(2)}. \tag{2.10}$$

Therefore

$$\Phi(t, z) = a\Phi^{(1)} + b\Phi^{(2)} \quad (2.11)$$

is the most general solution to the following problem

$$L\Phi = z\Phi \quad \text{and} \quad \frac{\partial \Phi}{\partial t_n} = (L^n)_+ \Phi, \quad (2.12)$$

with

$$\frac{\partial L}{\partial t_n} = [(L^n)_+, L]. \quad (2.13)$$

Lemma 2.1. *If $(\tau_n)_{n \in \mathbb{Z}}$ is a τ -vector for 1-Toda, then for arbitrary $z \in \mathbb{C}^*$, $a, b \in \mathbb{C}$, the vectors⁵*

$$((a\mathbb{X}_1(t, z) + be^{\xi(z)}\mathbb{X}_2(t, z))\tau)_{n \in \mathbb{Z}} \quad (2.14)$$

and

$$((1 + c\mathbb{X}(t, z))\tau)_{n \in \mathbb{Z}} \quad (2.15)$$

are 1-Toda vectors. The vertex operators $\mathbb{X}_1(y)$ and $e^{\xi(t, z)}\mathbb{X}_2(z)$ satisfy the commutation relation

$$\frac{y}{1 - y/z} \mathbb{X}_1(y) e^{\xi(z)} \mathbb{X}_2(z) = e^{\xi(z)} \mathbb{X}_2(z) \mathbb{X}_1(y). \quad (2.16)$$

Proof. According to the reduction above from 2- to 1-Toda, the tau-functions τ_n are independent of the sum of the arguments. Thus, we may write $\tau_n(x - y) = \tau_n(t)$ for $\tau_n(x, y)$. Set $\lambda = \mu^{-1} = z \in \mathbb{C}^*$ in (1.10); moreover, it is legitimate to multiply the τ -vector with an exponential in y_i with constant coefficients. Therefore the following expression is a 1-Toda τ -vector:

$$\begin{aligned} & e^{-\sum_1^\infty y_i z^i} \left((a\mathbb{Y}_1(z) + b\mathbb{Y}_2(z^{-1})) \tau \right)_n \\ &= e^{-\sum y_i z^i} \left(ae^{\sum_1^\infty x_i z^i} z^n \tau_n(x - y - [z^{-1}]) + be^{\sum y_i z^i} z^{-n} \tau_{n+1}(x - y + [z^{-1}]) \right) \\ &= ae^{\sum t_i z^i} z^n \tau_n(t - [z^{-1}]) + bz^{-n} \tau_{n+1}(t + [z^{-1}]) \\ &= ((a\mathbb{X}_1(z) + be^{\xi(z)}\mathbb{X}_2(z))\tau)_n. \end{aligned}$$

⁵ $\xi(t, z) := \sum_1^\infty t_i z^i$.

Finally, applying the inverse of $e^{\xi(y)}\mathbb{X}_2(y)$ to the above equation and taking a limit, when $y \rightarrow z$, leads to the result (2.15), after noting that

$$\lim_{y \rightarrow z} \left(e^{\xi(y)}\mathbb{X}_2(y) \right)^{-1} \left(e^{\xi(z)}\mathbb{X}_2(z) \right) = \lim_{y \rightarrow z} \Lambda^{-1} \chi(yz^{-1}) e^{\sum \frac{z^{-i}-y^{-i}}{i} \frac{\partial}{\partial t_i}} \Lambda = I$$

$$\begin{aligned} & \lim_{y \rightarrow z} \frac{1}{1 - z/y} \left(e^{\xi(y)}\mathbb{X}_2(y) \right)^{-1} \mathbb{X}_1(z) \\ &= \lim_{y \rightarrow z} \frac{1}{1 - z/y} \Lambda^{-1} \chi(y) \chi(z) e^{-\sum \frac{y^{-i}}{i} \frac{\partial}{\partial t_i}} e^{\sum t_i z^i} e^{-\sum \frac{z^{-i}}{i} \frac{\partial}{\partial t_i}} \\ &= \Lambda^{-1} \chi(z^2) e^{\sum t_i z^i} e^{-2 \sum \frac{z^{-i}}{i} \frac{\partial}{\partial t_i}} \\ &= \mathbb{X}(t, z), \end{aligned}$$

using the commutation relation

$$[e^{-\sum \frac{y^{-i}}{i} \frac{\partial}{\partial t_i}}, e^{\sum t_i z^i}] = -\frac{z}{y} e^{\sum t_i z^i} e^{-\sum \frac{y^{-i}}{i} \frac{\partial}{\partial t_i}}.$$

■

3 Toda-Darboux transformations

The purpose of this section is to establish theorem 0.1, which we restate in an explicit form.

Proposition 3.1. *For L evolving according to the Toda lattice equations (2.9), the Borel decomposition $L(t) - \lambda = L_-(t)L_+(t)$ is given for all $t \in \mathbb{C}^\infty$, by*

$$\alpha_n = \frac{\partial}{\partial t_1} \log \left(e^{-\sum t_i \lambda^i} \Phi_{n+1}(t, \lambda) \right) = \frac{\partial}{\partial t_1} \log \Phi_{n+1}(t, \lambda) - \lambda \quad (3.1)$$

$$\beta_n = -\frac{\partial}{\partial t_1} \log \left(\frac{\tau_n}{\tau_{n+1}} \Phi_n(t, \lambda) \right) = -\frac{\Phi_{n+1}(t, \lambda)}{\Phi_n(t, \lambda)} \quad (3.2)$$

in terms of the vector Φ , defined in (2.11), i.e.,

$$L - \lambda = L_- L_+ = \left(\Lambda^{-1} \left(\Lambda \frac{\partial}{\partial t_1} \log \Phi(t, \lambda) - \lambda \right) + \Lambda^0 \right) \left(- \left(\frac{\Lambda \Phi(t, \lambda)}{\Phi(t, \lambda)} \right) \Lambda^0 + \Lambda \right);$$

so, Φ belongs to the kernel of L_+ , i.e.,

$$L_+ \Phi = 0.$$

Finally the Darboux map

$$L(t) - \lambda = L_-(t) L_+(t) \mapsto \tilde{L}(t) - \lambda = L_+(t) L_-(t)$$

is Toda-Darboux, i.e., $\tilde{L}(t)$ satisfies the Toda lattice as well.

Proof: Let us begin by showing the second identity in (3.2), using (0.3) and (2.12):

$$\Phi_n \frac{\partial}{\partial t_1} \log \left(\frac{\tau_n}{\tau_{n+1}} \Phi_n(t, \lambda) \right) = -b_n \Phi_n + \frac{\partial \Phi_n}{\partial t_1} = -b_n \Phi_n + (L_+ \Phi)_n = \Phi_{n+1}.$$

To establish the theorem, we must check, at first, that $L - \lambda = L_- L_+$, with the entries α_n and β_n , given by (3.1) and (3.2); secondly, we must check that $\tilde{L} - \lambda = L_+ L_-$ evolves according to the Toda lattice.

To begin with, we verify the transformation (0.6): $b_n - \lambda = \alpha_{n-1} + \beta_n$ and $a_{n-1} = \alpha_{n-1} \beta_{n-1}$, with α_n, β_n , and b_n equal to (3.1), (3.2) and (0.3) respectively:

$$\alpha_{n-1} + \lambda + \beta_n - b_n = \frac{\partial}{\partial t_1} \left(\log \Phi_n(t, \lambda) - \log \frac{\tau_n}{\tau_{n+1}} \Phi_n(t, \lambda) - \log \frac{\tau_{n+1}}{\tau_n} \right) = 0$$

$$\begin{aligned} \Phi_{n-1}(a_{n-1} - \alpha_{n-1} \beta_{n-1}) &= \Phi_{n-1} \left(a_{n-1} + \left(\frac{\partial}{\partial t_1} \log \Phi_n - \lambda \right) \frac{\Phi_n}{\Phi_{n-1}} \right) \\ &= a_{n-1} \Phi_{n-1} + \frac{\partial \Phi_n}{\partial t_1} - \lambda \Phi_n \\ &= \left(\left(\frac{\partial}{\partial t_1} - L + L_- \right) \Phi \right)_n = 0. \end{aligned}$$

Remember from (0.6), that the Darboux map

$$L - \lambda = L_- L_+ \mapsto \tilde{L} - \lambda = L_+ L_- \quad (3.3)$$

reads componentwise:

$$b_n - \lambda = \alpha_{n-1} + \beta_n \longmapsto \tilde{b}_n - \lambda = \alpha_n + \beta_n = (b_n - \lambda) + (\alpha_n - \alpha_{n-1})$$

$$a_{n-1} = \alpha_{n-1}\beta_{n-1} \longmapsto \tilde{a}_{n-1} = \alpha_{n-1}\beta_n = a_{n-1}\frac{\beta_n}{\beta_{n-1}}$$

with α_n and β_n given by (3.2). So, in terms of τ_n and Φ_n , we have

$$\begin{aligned} \tilde{b}_n - \lambda &= (b_n - \lambda) + (\alpha_n - \alpha_{n-1}) \\ &= \frac{\partial}{\partial t_1} \log \frac{\tau_{n+1}}{\tau_n} - \lambda + \frac{\partial}{\partial t_1} \log \frac{\Phi_{n+1}(t, \lambda)}{\Phi_n(t, \lambda)} \\ &= \frac{\partial}{\partial t_1} \log \frac{\tau_{n+1}\Phi_{n+1}(t, \lambda)}{\tau_n\Phi_n(t, \lambda)} - \lambda \\ &= \frac{\partial}{\partial t_1} \log \frac{\tilde{\tau}_{n+1}}{\tilde{\tau}_n} - \lambda \\ \tilde{a}_{n-1} &= a_{n-1} \frac{\beta_n}{\beta_{n-1}} \\ &= \frac{\tau_{n-1}\tau_{n+1}}{\tau_n^2} \cdot \frac{\Phi_{n-1}(t, \lambda)\Phi_{n+1}(t, \lambda)}{\Phi_n(t, \lambda)^2} \\ &= \frac{\tilde{\tau}_{n-1}\tilde{\tau}_{n+1}}{\tilde{\tau}_n^2}, \end{aligned}$$

where

$$\tilde{\tau} = \tau(t)\Phi(t, \lambda) = (a\mathbb{X}_1(\lambda) + be^{\xi(\lambda)}\mathbb{X}_2(\lambda)) \tau$$

is a new τ -vector, by virtue of Lemma 2.1. Hence, the matrix $\tilde{L}$, parametrized by this new τ -vector $\tilde{\tau}$, satisfies the symmetric 1-Toda equations. ■

4 Wronskians and Vertex operators for 1-Toda

The aim of this section is to prove Theorem 0.2. The Wronskian on vectors $f = (f_n)_{n \in \mathbb{Z}}$ was defined in (0.9), the vertex operators $\mathbb{X}_1(t, y)$ and $\mathbb{X}_2(t, z)$ in (0.7), and $\xi(t, z) = \xi(z) = \sum_1^\infty t_i z^i$.

Proposition 4.1. *The following identities hold:*

$$\begin{aligned}\{\Phi^{(1)}(t, y), \Phi^{(1)}(t, z)\} &= y \frac{\mathbb{X}_1(y) \mathbb{X}_1(z) \tau}{\tau} \\ \{\Phi^{(1)}(t, y), \Phi^{(2)}(t, z)\} &= y \frac{\mathbb{X}_1(y) e^{\xi(z)} \mathbb{X}_2(z) \tau}{\tau} = \left(1 - \frac{y}{z}\right) \frac{e^{\xi(z)} \mathbb{X}_2(z) \mathbb{X}_1(y) \tau}{\tau} \\ \{\Phi^{(2)}(t, y), \Phi^{(2)}(t, z)\} &= - \left(1 - \frac{z}{y}\right) \frac{e^{\xi(y)} \mathbb{X}_2(y) e^{\xi(z)} \mathbb{X}_2(z) \tau}{\tau}.\end{aligned}$$

Proof. Using, in the third equality, Fay identity (1.9) with the shift $t \mapsto t - [z_1^{-1}]$, and the limits $z_1 \rightarrow z$, $z_2 \rightarrow y$ and $v_1 = v_2 \rightarrow 0$,

$$\begin{aligned}\{\Phi_n^{(1)}(t, y), \Phi_n^{(1)}(t, z)\} &= \frac{e^{\xi(y) + \xi(z)}}{\tau_n \tau_{n+1}} (yz)^n \\ &\quad (y \tau_{n+1}(t - [y^{-1}]) \tau_n(t - [z^{-1}]) - z \tau_n(t - [y^{-1}]) \tau_{n+1}(t - [z^{-1}])) \\ &= \frac{e^{\xi(y) + \xi(z)}}{\tau_n \tau_{n+1}} (yz)^n (y - z) \tau_{n+1} \tau_n (t - [y^{-1}] - [z^{-1}]) \\ &= e^{\xi(y) + \xi(z)} (y - z) (yz)^n \frac{\tau_n(t - [y^{-1}] - [z^{-1}])}{\tau_n(t)} \\ &= y \left(\frac{\mathbb{X}_1(y) \mathbb{X}_1(z) \tau}{\tau} \right)_n.\end{aligned}$$

Using in the third equality Fay identity (1.8) with $z \rightarrow y$, $u \rightarrow z^{-1}$, $v \rightarrow 0$ and $n \mapsto n + 1$:

$$\begin{aligned}\{\Phi_n^{(1)}(t, y), \Phi_n^{(2)}(t, z)\} &= \frac{e^{\xi(y)}}{\tau_n \tau_{n+1}} \left(\frac{y}{z}\right)^n \\ &\quad (y \tau_{n+1}(t - [y^{-1}]) \tau_{n+1}(t + [z^{-1}]) - z^{-1} \tau_n(t - [y^{-1}]) \tau_{n+2}(t + [z^{-1}])) \\ &= \frac{e^{\xi(y)}}{\tau_n \tau_{n+1}} \left(\frac{y}{z}\right)^n y \tau_{n+1}(t - [y^{-1}] + [z^{-1}]) \tau_{n+1} \\ &= e^{\xi(y)} y \left(\frac{y}{z}\right)^n \frac{\tau_{n+1}(t - [y^{-1}] + [z^{-1}])}{\tau_n} \\ &= y \left(\frac{\mathbb{X}_1(y) e^{\xi(z)} \mathbb{X}_2(z) \tau}{\tau} \right)_n = \left(1 - \frac{y}{z}\right) \left(\frac{e^{\xi(z)} \mathbb{X}_2(z) \mathbb{X}_1(y) \tau}{\tau} \right)_n,\end{aligned}$$

using the commutation relation (2.16), in the last equality.

Using in the third equality the same identity as for the first wronskian, but with $t \mapsto t + [y^{-1}] + [z^{-1}]$ and $n \mapsto n + 1$:

$$\begin{aligned}
 & \{\Phi_n^{(2)}(t, y), \Phi_n^{(2)}(t, z)\} \\
 &= \frac{1}{\tau_n \tau_{n+1}} (yz)^{-n-1} \\
 & \quad (z\tau_{n+2}(t + [y^{-1}])\tau_{n+1}(t + [z^{-1}]) - y\tau_{n+1}(t + [y^{-1}])\tau_{n+2}(t + [z^{-1}])) \\
 &= -\frac{1}{\tau_n \tau_{n+1}} (yz)^{-n-1} (y - z)\tau_{n+2}(t + [y^{-1}] + [z^{-1}])\tau_{n+1} \\
 &= -(y - z)(yz)^{-n-1} \frac{\tau_{n+2}(t + [y^{-1}] + [z^{-1}])}{\tau_n} \\
 &= -\left(1 - \frac{z}{y}\right) \left(\frac{e^{\xi(y)}\mathbb{X}_2(y)e^{\xi(z)}\mathbb{X}_2(z)\tau}{\tau}\right)_n
 \end{aligned}$$

■

Proof of Theorem 0.2: Proposition 4.1 implies the following relations:

$$\{\Phi_n^{(1)}(t, z), \Phi_n(t, \lambda)\} = z \left(\frac{\mathbb{X}_1(z)(a\mathbb{X}_1(\lambda) + be^{\xi(\lambda)}\mathbb{X}_2(\lambda))\tau}{\tau} \right)_n = z \frac{X_1(z)\tilde{\tau}}{\tau},$$

and

$$\begin{aligned}
 \{\Phi_n^{(2)}(t, z), \Phi_n(t, \lambda)\} &= -(1 - \frac{\lambda}{z}) \left(\frac{e^{\xi(z)}\mathbb{X}_2(z)(a\mathbb{X}_1(\lambda) + be^{\xi(\lambda)}\mathbb{X}_2(\lambda))\tau}{\tau} \right)_n \\
 &= \frac{\lambda - z}{z} \frac{e^{\xi(z)}X_2(z)\tilde{\tau}}{\tau},
 \end{aligned}$$

from which the proof follows.

■

5 Borel factorization, KM-lattice and Ricatti equations

This section concerns itself with the proof of Theorem 0.3. For a fixed $\lambda \in \mathbb{C}^*$, consider the Darboux map

$$L - \lambda = L_- L_+ \longmapsto \tilde{L} - \lambda = L_+ L_-, \quad (5.1)$$

which in coordinates reads

$$\begin{aligned} b_n - \lambda &= \alpha_{n-1} + \beta_n \mapsto \tilde{b}_n - \lambda = \alpha_n + \beta_n = (b_n - \lambda) + (\alpha_n - \alpha_{n-1}) \\ a_{n-1} &= \alpha_{n-1}\beta_{n-1} \mapsto \tilde{a}_{n-1} = \alpha_{n-1}\beta_n = a_{n-1} \frac{\beta_n}{\beta_{n-1}}. \end{aligned} \quad (5.2)$$

Remember the first Toda flow ($= \partial/\partial t_1 = \cdot$)

$$b'_n = a_n - a_{n-1}, \quad a'_n = a_n(b_{n+1} - b_n) \quad (5.3)$$

and the Kac-Moerbeke lattice, referred to with $\cdot$:

$$\dot{\alpha}_n = (\beta_{n+1} - \beta_n)\alpha_n, \quad \dot{\beta}_n = (\alpha_n - \alpha_{n-1})\beta_n. \quad (5.4)$$

Proposition 5.1. *Toda for L and $\tilde{L} \iff$ KM for L_- and L_+ .*

Proof: Assuming the KM-vector field on (α, β) , one computes

$$(b_n - \lambda) \cdot = \dot{\alpha}_{n-1} + \dot{\beta}_n = \alpha_n \beta_n - \alpha_{n-1} \beta_{n-1} = a_n - a_{n-1} = (b_n - \lambda)' \quad (5.5)$$

$$\dot{a}_n = \dot{\alpha}_n \beta_n + \alpha_n \dot{\beta}_n = \alpha_n \beta_n (\beta_{n+1} - \beta_n + \alpha_n - \alpha_{n-1}) = a_n (b_{n+1} - b_n) = a'_n \quad (5.6)$$

and

$$(\tilde{b}_n - \lambda) \cdot = \dot{\alpha}_n + \dot{\beta}_n = \alpha_n \beta_{n+1} - \alpha_{n-1} \beta_n = \tilde{a}_n - \tilde{a}_{n-1} = (\tilde{b}_n - \lambda)' \quad (5.7)$$

$$\dot{\tilde{a}}_n = \dot{\alpha}_n \beta_{n+1} + \alpha_n \dot{\beta}_{n+1} = \alpha_n \beta_{n+1} (\beta_{n+1} - \beta_n + \alpha_{n+1} - \alpha_n) = \tilde{a}_n (\tilde{b}_{n+1} - \tilde{b}_n) = \tilde{a}'_n. \quad (5.8)$$

Conversely, assuming Toda on L and $\tilde{L}$ leads to the following four equations on α and β ; the first two are (5.5) and (5.8), with a shift, the last two are (5.6) and (5.7):

$$\begin{cases} \alpha'_n + \beta'_{n+1} &= \alpha_{n+1} \beta_{n+1} - \alpha_n \beta_n \\ \alpha'_n \beta_{n+1} + \alpha_n \beta'_{n+1} &= \alpha_n \beta_{n+1} (\beta_{n+1} - \beta_n + \alpha_{n+1} - \alpha_n) \end{cases}$$

and

$$\begin{cases} \alpha'_n \beta_n + \alpha_n \beta'_n &= \alpha_n \beta_n (\beta_{n+1} - \beta_n + \alpha_n - \alpha_{n-1}) \\ \alpha'_n + \beta'_n &= \alpha_n \beta_{n+1} - \alpha_n \beta_n. \end{cases}$$

Solving the first system in α'_n and β'_{n+1} or the second in α'_n and β'_n leads to the KM-lattice equations (5.4); so, now we may replace the differentiation $\cdot$ by $'$. ■

The KM-lattice equations (3.7)

$$\alpha'_n = (\beta_{n+1} - \beta_n)\alpha_n \quad \text{and} \quad \beta'_n = (\alpha_n - \alpha_{n-1})\beta_n,$$

upon using $\beta_n = \frac{a_n}{\alpha_n}$, $\beta_{n+1} = b_{n+1} - \alpha_n - \lambda$ in the first equation and $\alpha_n = \frac{a_n}{\beta_n}$ and $\alpha_{n-1} = b_n - \lambda - \beta_n$ in the second equation, yield

$$\begin{aligned} \alpha'_n &= -\alpha_n^2 + (b_{n+1} - \lambda)\alpha_n - a_n \\ \beta'_n &= \beta_n^2 - (b_n - \lambda)\beta_n + a_n \end{aligned} \quad (\text{Ricatti equations}); \quad (5.9)$$

the transformations

$$\alpha_n = \frac{\partial}{\partial t_1} \log \gamma_n \quad \text{and} \quad \beta_n = -\frac{\partial}{\partial t_1} \log \varepsilon_n \quad (5.10)$$

yield the second order linear equations

$$\gamma''_n - (b_{n+1} - \lambda)\gamma'_n + a_n\gamma_n = 0 \quad (5.11)$$

$$\varepsilon''_n + (b_n - \lambda)\varepsilon'_n + a_n\varepsilon_n = 0. \quad (5.12)$$

Proposition 5.2. *The expressions*

$$\alpha_n = \frac{\partial}{\partial t_1} \log \gamma_n := \frac{\partial}{\partial t_1} \log \left(e^{-\sum t_i \lambda^i} \Phi_{n+1}(t, \lambda) \right) \quad (5.13)$$

$$\beta_n = -\frac{\partial}{\partial t_1} \log \varepsilon_n := -\frac{\partial}{\partial t_1} \log \left(\frac{\tau_n}{\tau_{n+1}} \Phi_n(t, \lambda) \right), \quad (5.14)$$

given by the 2-dimensional family $\Phi(t, z) = a\Phi^{(1)} + b\Phi^{(2)}$, provide the most general solution to the Ricatti equations (5.9).

Proof. Since the parametrization of α_n and β_n in Proposition 3.1 provide a Toda-Darboux transformation, then, by proposition 5.1, α_n and β_n satisfy the KM-Lattice, and hence provide a 2-dimensional solution to the Ricatti equations (5.9).

An *alternate proof* not using τ -function theory proceeds as follows: The Ricatti equations (5.9) are equivalent via the transformation (5.2) and the Toda vector fields (5.3). We check, for instance, that ε_n satisfies (5.12); at first, we compute:

$$\Phi'_n = (L_+ \Phi)_n = b_n \Phi_n + \Phi_{n+1} = \lambda \Phi_n - a_{n-1} \Phi_{n-1}, \quad \text{using } L\Phi = \lambda\Phi. \quad (5.15)$$

Using the first expression for Φ'_n and Φ'_{n-1} and the Toda lattice equation for a'_{n-1} , we then find

$$\begin{aligned}\Phi''_n &= \lambda\Phi'_n - a'_{n-1}\Phi_{n-1} - a_{n-1}\Phi'_{n-1} \\ &= \lambda(b_n\Phi_n + \Phi_{n+1}) - a_{n-1}(b_n - b_{n-1})\Phi_{n-1} - a_{n-1}(b_{n-1}\Phi_{n-1} + \Phi_n) \\ &= (b_n + \lambda)\Phi_{n+1} + (b_n^2 - a_{n-1})\Phi_n, \text{ using } \lambda\Phi_n = (L\Phi)_n.\end{aligned}\quad (5.16)$$

Then, using

$$\frac{\tau_{n+1}}{\tau_n} \left(\frac{\tau_n}{\tau_{n+1}} \right)' = -b_n \quad \text{and} \quad \frac{\tau_{n+1}}{\tau_n} \left(\frac{\tau_n}{\tau_{n+1}} \right)'' = -b'_n + b_n^2 = a_{n-1} - a_n + b_n^2, \quad (5.17)$$

we find

$$\begin{aligned}\frac{\tau_{n+1}}{\tau_n}(\varepsilon''_n + (b_n - \lambda)\varepsilon'_n + a_n\varepsilon_n) \\ &= \Phi''_n - 2b_n\Phi'_n + (-b'_n + b_n^2)\Phi_n + (b_n - \lambda)(\Phi'_n - b_n\Phi_n) + a_n\Phi_n \\ &\stackrel{*}{=} \Phi''_n - (b_n + \lambda)\Phi_{n+1} + (-b_n^2 + a_{n-1})\Phi_n \\ &= 0, \quad \text{by (5.18)}\end{aligned}$$

in $\stackrel{*}{=}$, we have used (5.3), (5.17) and $\lambda\Phi = L\Phi$. ■

6 Toda flows and Toda-Darboux transforms for band matrices

In this section, we factorize band matrices of the form $L = \sum_{-p \leq i \leq p} a_i \Lambda^i$, and study their Toda-Darboux transforms.

Lemma 6.1. *Consider the difference operator*

$$L = a_{-r}\Lambda^{-r} + a_{-r+1}\Lambda^{-r+1} + \dots + a_{n-r-1}\Lambda^{n-r-1} + \Lambda^{n-r} \quad (6.1)$$

with $n \geq 2$, $r \geq 0$, diagonal matrices a_j , with leading term $a_{-r}(j) \neq 0$ for j sufficiently small. Then any choice of basis $\Phi^{(1)}, \dots, \Phi^{(n)} \in \ker L$ leads to a

factorization of⁶ L :

$$\begin{aligned} L = & (I - \Lambda^{-1}(\Lambda^{-r+1}\beta_n)) (I - \Lambda^{-1}(\Lambda^{-r+2}\beta_{n-1})) \dots (I - \Lambda^{-1}(\beta_{n-r+1})) \cdot \\ & \cdot (\Lambda - \beta_{n-r}I) (\Lambda - \beta_{n-r-1}I) \dots (\Lambda - \beta_1I), \end{aligned} \quad (6.2)$$

with

$$\beta_k = \frac{\Lambda\alpha_k}{\alpha_k}, \quad \alpha_k(\ell) = \frac{W_k(\ell)}{W_{k-1}(\ell)}, \quad W_k(\ell) = W[\Phi^{(1)}, \Phi^{(2)}, \dots, \Phi^{(k)}](\ell).$$

Proof: Step 1: $r = 0$

First we prove the statement for $r = 0$ by induction on the degree of L . Define the operator L_i ,

$$L_i(f) := \frac{W[f, \Phi^{(1)}, \dots, \Phi^{(i)}]}{W[\Phi^{(1)}, \dots, \Phi^{(i)}]}$$

which is the unique operator of the form (6.1) with $r = 0$, $n = i$ such that $\ker L_i = \{\Phi^{(1)}, \dots, \Phi^{(i)}\}$. But clearly

$$L_{i+1}(f) = \frac{W[L_i(f), L_i(\Phi^{(i+1)})]}{L_i(\Phi^{(i+1)})} = \left(\Lambda - \frac{\Lambda\alpha_i}{\alpha_i}I \right) L_i(f),$$

with $\alpha_i = L_i(\Phi^{(i+1)}) = W_{i+1}/W_i$. So, by induction $L_i(f)$ factors, thus leading to (6.2) for $r = 0$.

Step 2: $r \neq 0$

The case $r \neq 0$ is taken care of by multiplying (6.2) to the left with Λ^r :

$$\Lambda^r L = (\Lambda - \beta_n I) \dots (\Lambda - \beta_1 I),$$

on which you apply step 1. ■

In the next Lemma, we use definition (0.19) for $\hat{x}, \hat{y}, \hat{t}$; also define

$$\hat{x} - [\lambda^{-1}] := \left(x_i - \frac{\lambda^{-i}}{i} \right)_{i \neq p}, \quad \hat{t} - [\lambda^{-1}] = \left(t_{ip} - \frac{\lambda^{-ip}}{ip} \right)_{i=1,2,\dots}.$$

⁶Wronskian $= W_k(\ell) = W[\Phi^{(1)}, \Phi^{(2)}, \dots, \Phi^{(k)}](\ell) = \det \left(\Phi_{i+\ell-1}^{(j)} \right)_{1 \leq i, j \leq k}$. Note Λ^{-1} in the formula (6.2) refers to the expression in brackets only!

Lemma 2. *On the locus $L := L_1^p = L_2^p$, the 2-Toda vector fields (2.1) on (L_1, L_2) take the form (0.16), with τ -functions of the form (0.18) and with the $2p$ -dimensional null-space $\ker(L - \lambda^p)$ given by*

$$\Phi(\hat{x}, \hat{y}, \hat{t}; \lambda) = \sum_{0 \leq i \leq p-1} a_i \Phi^{(1)}(\omega^i \lambda) + \sum_{0 \leq i \leq p-1} b_i \Phi^{(2)}(\omega^i \lambda) \quad (6.3)$$

flowing according to the equations (0.20); in (6.3)

$$\begin{aligned} \Phi^{(1)}(\hat{x}, \hat{y}, \hat{t}; \lambda) &:= \Psi_1(x, y; \lambda) e^{-\sum_1^\infty y_{ip} \lambda^{ip}} \\ &= e^{\sum_{i \neq p} x_i \lambda^i} e^{\sum_{i=1}^\infty t_{ip} \lambda^{ip}} \left(z^n \frac{\tau_n(\hat{x} - [\lambda^{-1}], \hat{y}, \hat{t} - [\lambda^{-1}])}{\tau_n(\hat{x}, \hat{y}, \hat{t})} \right)_{n \in \mathbb{Z}} \\ &= \frac{\mathbb{X}_1(\lambda) \tau}{\tau} \end{aligned}$$

$$\begin{aligned} \Phi^{(2)}(\hat{x}, \hat{y}, \hat{t}; \lambda) &:= \Psi_2(x, y; \lambda^{-1}) e^{-\sum_1^\infty y_{ip} \lambda^{ip}} \\ &= e^{\sum_{i \neq p} y_i \lambda^i} \left(z^{-n} \frac{\tau_{n+1}(\hat{x}, \hat{y} - [\lambda^{-1}], \hat{t} + [\lambda^{-1}])}{\tau_n(\hat{x}, \hat{y}, \hat{t})} \right)_{n \in \mathbb{Z}} \\ &= \frac{e^{\sum_1^\infty t_{ip} \lambda^{ip}} \mathbb{X}_2(\lambda) \tau}{\tau}. \end{aligned}$$

Proof: The proof proceeds as in section 2; for instance, (2.2) gets replaced by

$$\frac{\partial(L_1^p - L_2^p)}{\partial x_n} = 0, \quad \frac{\partial(L_1^p - L_2^p)}{\partial y_n} = 0$$

and (2.3) by

$$\left(\frac{\partial}{\partial x_{np}} + \frac{\partial}{\partial y_{np}} \right) L_i = [(L_1^{np})_+ + (L_2^{np})_-, L_i] = [(L_i^{np})_+ + (L^{np})_-, L_i] = 0.$$

That Φ satisfies the differential equations (0.20) proceeds along the same lines as (2.9) and (2.10). $\blacksquare$

Proof of Theorem 0.4: Lemma 2 implies Theorem 0.4, except for the statement about the Darboux transform. For that, one needs to factor L according to the recipe of Lemma 1, where we set $\Phi = \Phi^{(1)}$, in the precise notation of Lemma 1. Thus the Darboux transformation corresponds to bringing the factor most to the right all the way to the left. $\blacksquare$

References

- [1] M. Adler: *On the Bäcklund Transformation for the Gel'fand-Dickey Equations*, Commun. Math. Phys. 80, 517–527 (1981).
- [2] M. Adler and P. van Moerbeke: *Birkhoff strata, Bäcklund transformations and limits of isospectral operators*, Adv. in Math., **108** 140–204 (1994).
- [3] M. Adler and P. van Moerbeke: *Matrix integrals, Toda symmetries, Virasoro constraints and orthogonal polynomials* Duke Math.J., **80** (3), 863–911 (1995) and *String orthogonal Polynomials, String Equations and two-Toda Symmetries*, Comm. Pure and Appl. Math., **L** 241–290 (1997).
- [4] M. Adler and P. van Moerbeke: *Finite random Hermitean ensembles and The spectrum of coupled random matrices*, to appear (1997).
- [5] M. Adler, T. Shiota and P. van Moerbeke : *From the w_∞ -algebra to its central extension: a τ -function approach*, Physics Letters **A 194**, 33–43 (1994) and *A Lax pair representation for the Vertex operator and the central extension*, Commun. Math. Phys. 171, 547–588 (1995).
- [6] B. Bakalov, E. Horozov and M. Yakimov: *Bäcklund-Darboux transformations in Sato's Grassmannian*, reprint, 1996.
- [7] E. Date, M. Jimbo, M. Kashiwara, T. Miwa: *Transformation groups for soliton equations*, In: Proc. RIMS Symp. Nonlinear integrable systems — Classical and quantum theory (Kyoto 1981), pp. 39–119. Singapore : World Scientific 1983.
- [8] J. Fastré: *Boson-correspondence for W -algebras, Bäcklund-Darboux transformations and the equation $[L, P] = L^n$* , Bull. des Sciences Math. (1997)
- [9] V. B. Matveev, M.A. Salle: *Darboux transformations and solitons*, Springer series in Non-linear Dynamics, 1991.
- [10] K. Takasaki, T. Takebe: *Integrable hierarchies and dispersionless limit*, Reviews in Math. Phys. **7**, 743–808 (1995).

- [11] K. Ueno and K. Takasaki: *Toda Lattice Hierarchy*, Adv. Studies in Pure Math. **4**, 1–95 (1984).
- [12] P. van Moerbeke : *Integrable foundations of string theory*, in Lectures on Integrable systems, Proceedings of the CIMPA-school, 1991, Ed.: O. Babelon, P. Cartier, Y. Kosmann-Schwarzbach, World scientific, 163–267 (1994).